

03105-38-1

THE UNIVERSITY OF MICHIGAN

COLLEGE OF LITERATURE, SCIENCE, AND THE ARTS
DEPARTMENT OF COMMUNICATION SCIENCES

Technical Report

Transparent Categories and Categories of Transition Systems

YEHOSEFAT GIVE'ON

AD 466445

DDC

JUL 22 1965

1965 B

Under Contract with:

Department of the Navy
Office of Naval Research
Contract No. Nonr-1224(21)
Washington, D. C.

Administered through:

May 1965

OFFICE OF RESEARCH ADMINISTRATION • ANN ARBOR

BEST AVAILABLE COPY

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

THE UNIVERSITY OF MICHIGAN
COLLEGE OF LITERATURE, SCIENCE, AND THE ARTS
Department of Communication Sciences

Technical Report

TRANSPARENT CATEGORIES AND CATEGORIES OF TRANSITION SYSTEMS

Yehoshafat Give'on

ORA Project 03105

under contract with:

DEPARTMENT OF THE NAVY
OFFICE OF NAVAL RESEARCH
CONTRACT NO. Nonr-1224(21)
WASHINGTON, D.C.

administered through:

OFFICE OF RESEARCH ADMINISTRATION

ANN ARBOR

May 1965

RESEARCH PROGRESS REPORT

Title: "Transparent Categories and Categories of Transition Systems"
Y. Give'on, University of Michigan Technical Report 03105-38-T, May 1965;
Nonr 1224(21).

Background: The Logic of Computers Group of the Communication Sciences Department of The University of Michigan is investigating the application of logic and mathematics to the design of computing automata. The application of the techniques and concepts of abstract algebra to automata forms a part of this investigation.

Condensed Report Contents: Several recent results in automata theory give evidence of the importance of homomorphisms in the study of transition systems and automata. It is natural therefore to inquire how much information can be retrieved from the algebra of homomorphism compositions with respect to transition systems. The natural mathematical framework for the discussion of this problem is categorical algebra.

We define a category \mathbf{A}_W of the transition systems with input W , where W is any arbitrary fixed monoid, and with arbitrary sets of states. A preliminary study of \mathbf{A}_W (Give'on 1964) shows that one can reconstruct the internal structure of any transition system from the way homomorphisms (i.e., the morphisms of \mathbf{A}_W) behave around it.

In this paper we show that \mathbf{A}_W has a generator, M_W (which is W operating on itself as a transition system) and that there exists a functor $\text{Mor} : \mathbf{A}_W \rightarrow \mathbf{A}_W$ naturally equivalent to the identity functor of \mathbf{A}_W which factors through $\text{Hom}_{\mathbf{A}_W}(M_W, -)$.

A general exposition of the nature of properties which are retrievable from the "morphism-behavior" in an arbitrary category is presented so that it provides a rigorous general basis for studying "retrievable" properties and categories in which every structural property of objects and morphisms is "retrievable."

Finally, we prove that for a very broad class of input monoids, which includes all the types of input-monoids encountered in automata theory, the categories \mathbf{A}_W are transparent. That is, anything which can be said about the structure of transition systems with input W , can be said by referring to their homomorphisms only. In particular, all the automorphisms of \mathbf{A}_W , for this type of W , are naturally equivalent to the identity functor of \mathbf{A}_W .

For further information: The complete report is available in the major Navy technical libraries and can be obtained from the Defense Documentation Center. A few copies are available for distribution by the author.

1. INTRODUCTION

Several recent results in automata theory (in particular, Hartmanis & Stearns 1964, Zeiger 1964) give evidence of the importance of homomorphisms in the study of transition systems and automata. It is natural therefore to inquire how much information can be retrieved from the algebra of homomorphism compositions with respect to transition systems. The natural mathematical framework for the discussion of this problem is categorical algebra.

We define a category \mathbf{A}_W of the transition systems with input W , where W is any arbitrary fixed monoid, and with arbitrary sets of states. A preliminary study of \mathbf{A}_W (Give'on 1964) shows that one can reconstruct the internal structure of any transition system from the way homomorphisms (i.e., the morphisms of \mathbf{A}_W) behave around it.

In this paper we show that \mathbf{A}_W has a generator, M_W (which is W operating on itself as a transition system) and that there exists a functor $\text{Mor} : \mathbf{A}_W \rightarrow \mathbf{A}_W$ naturally equivalent to the identity functor of \mathbf{A}_W which factors through $\text{Hom}_{\mathbf{A}_W}(M_W, -)$.

A general exposition of the nature of properties which are retrievable from the "morphism-behavior" in an arbitrary category is presented so that it provides a rigorous general basis for studying "retrievable" properties and categories in which every structural property of objects and morphisms is "retrievable."

Finally, we prove that for a very broad class of input monoids, which includes all the types of input-monoids encountered in automata theory, the categories \mathcal{A}_W are transparent. That is, anything which can be said about the structure of transition systems with input W , can be said by referring to their homomorphisms only. In particular, all the automorphisms of \mathcal{A}_W , for this type of W , are naturally equivalent to the identity functor of \mathcal{A}_W .

Some elementary acquaintance with categorical algebra is needed. In particular, we shall make use of the following notions:

- (i) Category, its objects and its morphisms.
- (ii) Epic, monic, and invertible morphisms versus surjective, injective, and bijective functions.

(iii) Initial and terminal objects.

(iv) Functors, natural transformations and natural equivalences
of functors.

(v) Embedding functors, automorphism functors, and adjoint
functors.

The reader who is not familiar with these notions is referred
to the literature (Kan 1958, Freyd 1964, and MacLane 1965). Additional
issues of categorical algebra with reference to automata theory are discussed
in (Giv'e'on 1965).

2. CATEGORIES OF TRANSITION SYSTEMS

2.1 Let W be a fixed monoid. We denote by \mathcal{A}_W the category specified as follows.

The objects of \mathcal{A}_W are transition-systems with input W . That is, systems of the form

$$A = (S(A) \times W \xrightarrow{\lambda_A} S(A))$$

where:

(i) $S(A)$ is any set, the set of states of A ;

(ii) $\lambda_A : S(A) \times W \rightarrow S(A)$ is a function, the transition function of A , with the following properties (we write $s \cdot \omega$ for $\lambda_A(s, \omega)$):

(iii) $s \cdot 1_W = s$ for all $s \in S(A)$, where 1_W is the identity element of W ;

(iv) $s \cdot (\omega_1 \omega_2) = (s \cdot \omega_1) \cdot \omega_2$ for all $s \in S(A)$ and all $\omega_1, \omega_2 \in W$.

The morphisms of \mathcal{A}_W are of the form

$$A \xrightarrow{f} B$$

where $f : S(A) \rightarrow S(B)$ is a function satisfying $f(s \cdot \omega) = f(s) \cdot \omega$ for all $s \in S(A)$ and all $\omega \in W$. (Note that $s \cdot \omega$ on the left hand of this equation refers to the transition function of A , while $f(s) \cdot \omega$ refers to the transition function of B .)

The composition of the morphisms of \mathbf{A}_W is determined in an obvious manner by the composition of the functions which underly the morphisms. That is, $(C \xrightarrow{g} D) (A \xrightarrow{f} B)$ is defined only when $B = C$ and then it is equal to $A \xrightarrow{gf} D$.

2.2 As in many other "natural" categories of mathematical systems, we have a forgetful functor $S : \mathbf{A}_W \rightarrow \mathbf{S}$ from \mathbf{A}_W to \mathbf{S} the category of sets where $S(A)$ is the set of states of A and $S(A \xrightarrow{f} B) = (f : S(A) \rightarrow S(B))$.

Note that \mathbf{A}_W contains, among its objects, an empty object to be denoted by ϕ_A . Here we adopt the useful convention that for any set T there exists a unique function which is injective (i.e., one-one into) from ϕ , the empty set, into T . Thus, the transition function of ϕ_W is this "empty" function : $\phi \times W \rightarrow \phi$ ($\phi \times W = \phi$) and for any object A of \mathbf{A}_W there exists a unique morphism $\phi_W \xrightarrow{\sigma_A} A$ which is determined by the "empty"

$$\phi = S(\phi_W) \longrightarrow S(A).$$

The forgetful functor $S : \mathcal{A}_W \longrightarrow S$ has an adjoint (cf. Kan 1958, MacLane 1965), the functor $Fr : S \longrightarrow \mathcal{A}_W$, which assigns to each set T , an object $Fr(T)$ of \mathcal{A}_W which is free on $T \subseteq S(Fr(T))$.

The functor $Fr : S \longrightarrow \mathcal{A}_W$ can be specified as follows. For any set T , $Fr(T)$ is the transition system defined by:

$$S(Fr(T)) = T \times W,$$

$$(t, \omega_1) \cdot \omega_2 = (t, \omega_1 \omega_2).$$

For any function $f : T_1 \longrightarrow T_2$ there exists a unique morphism

$$Fr(T_1) \xrightarrow{Fr(f)} Fr(T_2) \text{ such that for any } t \in T_1 : [Fr(f)](t, 1_W) = (f(t), 1_W).$$

Hence $Fr(T)$ is "free on $T \times \{1_W\}$ ". We identify the elements of $T \times \{1_W\}$

with the elements of T : $t \equiv (t, 1_W)$.

If T_1 and T_2 are sets which have the same cardinality, then

$Fr(T_1)$ and $Fr(T_2)$ are isomorphic (i.e., there exists an invertible

morphism $Fr(T_1) \longrightarrow Fr(T_2)$ of \mathcal{A}_W . In particular, if T is a single-element

set then we denote $Fr(T)$ by M_W .

M_W serves a very important role in A_W as we shall see later.

Note that M_W may be defined as W operating on itself. That is,

$$S(M_W) = W,$$

$$\omega_1 \cdot \omega_2 = \omega_1 \omega_2 \quad .$$

2.3 A_W shares with the obelian "natural" categories, e.g., of groups or of modules, (cf. Fröyd 1964) the property that the monic (respectively, the epic, and the invertible) morphisms are precisely those morphisms of A_W whose underlying functions are injective (respectively, surjective and bijective). The arguments that establish these facts are similar to the arguments employed in the category of groups for the same end.

The existence of the forgetful functor $S : A_W \longrightarrow S$ implies that a morphism $A \xrightarrow{f} B$ is invertible in A_W iff f is bijective. Since S is an embedding functor, every morphism of A_W whose underlying function is injective (respectively, surjective) must be monic (respectively, epic).

In order to prove the converse (for monic and epic morphisms of A_W) we need some additional observations about A_W . These observations

will be incorporated in the proofs of the following lemmata.

2.3.1 LEMMA: If $A \xrightarrow{e} B$ is an epic morphism of \mathcal{A}_W then $e : S(A) \rightarrow S(B)$ is surjective.

Proof: The image of $e : S(A) \rightarrow S(B)$ is a subset $e(S(A))$ of $S(B)$ such that for any $\omega \in W$ and any $s \in e(S(A))$, $s \cdot \omega \in e(S(A))$. Hence $e(S(A))$ is a transition system $e(A)$ which is a sub-system of B .

We define a new object $B/e(A)$ of \mathcal{A}_W by:

$$S(B/e(A)) = (S(B) - e(S(A))) \cup \{s_*\} \text{ where } s_* \in S(B);$$

the transition function of $B/e(A)$ is the same as of B except for the cases where $s \cdot \omega \in e(S(A))$; in these cases we set $s \cdot \omega = s_*$, and for all $\omega \in W$ we set $s_* \cdot \omega = s_*$.

Obviously, $B/e(A)$ is formed from B by contracting $e(S(A))$ to a single state s_* . This contraction takes the form of a canonical morphism $B \xrightarrow{q_e} B/e(A)$, where $q_e : S(B) \rightarrow S(B/e(A))$ is identical on $S(B) - e(S(A))$ and it maps all of $e(S(A))$ onto s_* .

In addition to $B \xrightarrow{q_e} B/e(A)$, we have another morphism

$B \xrightarrow{z} B/e(A)$ which maps all of $e(S(A))$ onto s_* . Clearly e is surjective iff $q_e = z$.

Obviously $ze = q_e e$, since both map all of $S(A)$ onto s_* . But e is epic and therefore $ze = q_e e$ implies $q_e = z$.

2.3.2 LEMMA: If $A \xrightarrow{j} B$ is a monic morphism of \mathcal{A}_W then $j : S(A) \rightarrow S(B)$ is injective.

Proof: Assume that for $s_1, s_2 \in S(A)$ we have $j(s_1) = j(s_2)$.

We define two morphisms $M_W \xrightarrow{f_1} A$ and $M_W \xrightarrow{f_2} A$ by $f_1(l_W) = s_1$, and $f_2(l_W) = s_2$. Obviously, $j f_1 = j f_2$, and since j is monic, it follows that $f_1 = f_2$; i.e., $s_1 = s_2$.

2.4 For any object A of \mathcal{A}_W any any subset $T \subseteq S(A)$, we define $A(T)$,

the subsystem of A generated by T, as follows:

$$S(A(T)) = T \cdot W = \{t \cdot \omega : t \in T \text{ and } \omega \in W\},$$

$$(t \cdot \omega_1) \cdot \omega_2 = t \cdot (\omega_1 \omega_2).$$

A subset T of S(A) is said to generate A iff $A(T) = A$;

i.e., iff $T \cdot W = S(A)$. In particular A is said to be monogenic iff A

is generated by a single-element subset of S(A).

For example, M_W is monogenic since $\{1_W\}$ generates M_W (Obviously for any $\omega \in W : 1_W \cdot \omega = \omega$). More generally, M_W is generated by $\{u\}$ iff there exists $v \in W$ such that $uv = 1_W$.

Note that an object A of \mathcal{A}_W is monogenic iff for any $T \subseteq S(A)$ which generates A there exists $t \in T$ such that $\{t\}$ generates A.

2.4.1 LEMMA: An object A of \mathcal{A}_W is monogenic iff for any family

$\{A_j\}$ of subsystems of A indexed by a set J, $U\{S(A_j) : j \in J\} = S(A)$

implies $S(A_j) = S(A)$ for some $j \in J$.

Proof: Assume that A is monogenic and generated by $\{s_0\}$.

If $US(A_j) = S(A)$ then $s_0 \in S(A_j)$ for some $j \in J$, and $S(A_j) = S(A)$.

Assume that for any family $\{A_j\}$ indexed by a set J ,

$US(A_j) = S(A)$ implies $S(A_j) = S(A)$ for some $j \in J$. Define the family

$\{A_s\}$ for all $s \in S(A)$, where A_s is the subsystem of A generated by $\{s\}$.

Obviously $US(A_s) = S(A)$ and therefore there exists $s_0 \in S(A)$ for

which $S(A_{s_0}) = S(A)$. Hence A is generated by $\{s_0\}$.

2.4.2 COROLLARY: For any monogenic object A of \mathcal{A}_W and any automorphism

$F : \mathcal{A}_W \rightarrow \mathcal{A}_W$ of \mathcal{A}_W , $F(A)$ is also a monogenic object of \mathcal{A}_W .

Proof: We recall that an automorphism F of \mathcal{A}_W is a functor

$F : \mathcal{A}_W \rightarrow \mathcal{A}_W$ for which there exists a functor $G : \mathcal{A}_W \rightarrow \mathcal{A}_W$ such that

both $F \circ G$ and $G \circ F$ are equal to the identity functor of \mathcal{A}_W .

The families of subsystems of A are represented faithfully

by the families of monic morphisms of \mathcal{A}_W with range A . Given a set

J of monic morphisms $A_j \xrightarrow{j} A$ we define a category \mathcal{T} (which is a

subcategory of \mathcal{A}_W) whose objects are all the monic morphisms

$B \xrightarrow{b} A$ such that for any $j \in J$ there exists a monic $A_j \xrightarrow{b_j} B$ with $bb_j = j$. The morphisms of \mathcal{J} are of the form

$$(B_1 \xrightarrow{b_1} A) \xrightarrow{f} (B_2 \xrightarrow{b_2} A)$$

where $B_1 \xrightarrow{f} B_2$ is a morphism of \mathcal{A}_W with $B_2 f = b_1$.

For any set J of monic morphisms of \mathcal{A}_W with range A , the category \mathcal{J} has an initial object $U(J)$, which is unique up to an isomorphism of \mathcal{J} (which is an equivalence of monic morphisms in \mathcal{A}_W (cf. Freyd 1964, MacLane 1965)). $U(J)$ is a monic morphism of \mathcal{A}_W with range A and whose image is precisely the union of the images of the morphisms in J .

We can rephrase now Lemma 2.4 1 : An object A of \mathcal{A}_W is monogenic iff for any set J of monic morphisms of \mathcal{A}_W with range A , if $U(J)$ is an invertible morphism of \mathcal{A}_W (i.e., an isomorphism) then there is a $j \in J$ which is invertible.

Since this characterization of the monogenic objects in \mathcal{A}_W is preserved under the automorphisms of \mathcal{A}_W the proof follows.

2.5 In the proof of Cor. 2.4.2 we have shown that the property of being a monogenic object of \mathcal{A}_W , which was defined originally by "looking inside A," is in fact definable by means of general properties of morphisms in categories. Knowing the way morphisms behave around an object A is sufficient in order to determine whether A contains a state from which all the rest of the states of A are accessible. In other words, the property of being a monogenic object in \mathcal{A}_W is categorical. In Chapter 4 we shall present a rigorous explication of this notion. The properties of \mathcal{M}_W , that we shall derive in the next chapter, will yield the result that all properties of objects of \mathcal{A}_W (which are invariant under isomorphisms in \mathcal{A}_W) are categorical (provided that W belong to a very broad class of monoids). That is, if W satisfies some weak conditions, then all the properties of the transition systems with input W can be derived from the categorical-algebra study of \mathcal{A}_W .

3. A STUDY OF M_W

3.1 LEMMA: An object A of \mathcal{A}_W is monogenic iff there exists an epic morphism $M_W \xrightarrow{e} A$.

Proof: Assume that A is monogenic and generated by $\{s_0\}$.

Define a morphism $M_W \xrightarrow{f_{s_0}} A$ by $f_{s_0}(1_W) = s_0$ (recall that M_W is free on 1_W). Obviously, $f_{s_0} : W \rightarrow S(A)$ is surjective and therefore $M_W \xrightarrow{f_{s_0}} A$ is epic.

On the other hand, if $M_W \xrightarrow{e} A$ is epic then A is generated by $e(1_W)$ since $e(\omega) = e(1_W) \cdot \omega$.

3.2 We define a functor $H_{M_W} : \mathcal{A}_W \rightarrow S$ by:

$$H_{M_W}(A) = \text{Hom}_{\mathcal{A}_W}(M_W, A)$$

$$H_{M_W}(A \xrightarrow{f} B) = (\ell_f : \text{Hom}_{\mathcal{A}_W}(M_W, A) \rightarrow \text{Hom}_{\mathcal{A}_W}(M_W, B))$$

where $\ell_f(M_W \xrightarrow{g} A) = (M_W \xrightarrow{fg} B)$.

We define a transformation of functors $\varphi: S \rightarrow H_{M_W}$ as

follows:

For any object A of A_W , $\varphi(A): S(A) \rightarrow H_{M_W}(A)$ is given by

$$[\varphi(A)](s): W \rightarrow S(A): \omega \rightarrow s \cdot \omega.$$

In other words, $[\varphi(A)](s)$ is the morphism $M_W \xrightarrow{f_s} A$ which is determined by $f_s(1_W) = s$.

The function $\varphi(A): S(A) \rightarrow H_{M_W}(A)$ is bijective. It

is injective since $f_{s_1} = f_{s_2}$ implies $f_{s_1}(1_W) = f_{s_2}(1_W)$. It is sur-

jective since for any morphism $M_W \xrightarrow{g} A$ we have $f_{g(1_W)} = g$.

Furthermore, for any morphism $A \xrightarrow{g} B$ of A_W , and for any $s \in S(A)$

we have

$$g \circ f_s = f_{g(s)},$$

where $f_{g(s)} = [\varphi(B)](g(s))$.

For we clearly have

$$(g \circ f_s)(\omega) = g(s \cdot \omega) = g(s) \cdot \omega = f_{g(s)}(\omega).$$

From this follows directly, that for any morphism $A \xrightarrow{g} B$

of \mathcal{A}_W , the following diagram is commutative.

$$\begin{array}{ccc}
 S(A) & \xrightarrow{g} & S(B) \\
 \downarrow \rho(A) & & \downarrow \rho(B) \\
 H_{M_W}(A) & \xrightarrow{H_{M_W}(g)} & H_{M_W}(B)
 \end{array}$$

Thus we have proved:

3.2.1 PROPOSITION: The transformation $\rho: S \rightarrow H_{M_W}$ is a natural equivalence of functors.

3.2.2 The pair (M_W, ρ^{-1}) is a representation of the forgetful functor $S: \mathcal{A}_W \rightarrow \mathcal{S}$ (cf. MacLane 1965).

3.2.3. Since $S: \mathcal{A}_W \rightarrow \mathcal{S}$ is an embedding functor (i.e., one-one on the morphisms) it follows that H_{M_W} is also an embedding and therefore

M_W is a generator of \mathcal{A}_W (cf. Freyd 1964).

3.2.4 COROLLARY: M_W is a projective object of \mathcal{A}_W . (An object P of

a category \mathcal{C} is projective iff for any morphism $P \xrightarrow{g} B$ and any epic morphism $A \xrightarrow{e} B$ of \mathcal{C} there exists a morphism $P \xrightarrow{f} A$ of \mathcal{C} for which the following diagram is commutative:

$$\begin{array}{ccc} & P & \\ f \swarrow & \downarrow g & \\ A & \xrightarrow{e} & B \end{array} \quad .)$$

Proof: It is sufficient (and necessary) to show that if $A \xrightarrow{e} B$ is an epic morphism of \mathcal{A}_W then $H_{M_W}(e) : H_{M_W}(A) \rightarrow H_{M_W}(B)$ is surjective. From the commutative diagram for $\rho : S \rightarrow H_{M_W}$ we derive

$$H_{M_W}(e) = \rho(B)e(\rho(A))^{-1} \quad . \quad \text{Hence } H_{M_W}(e) \text{ is surjective.}$$

3.3.3 PROPOSITION: The bijection $\rho(M_W) : W \rightarrow H_{M_W}(M_W)$ determines

an isomorphism of monoids

$$W \xrightarrow{R} \text{End}_{\mathcal{A}_W}(M_W)$$

where $\text{End}_{\mathcal{A}_W}(M_W)$ is the monoid of the morphisms $M_W \rightarrow M_W$ of \mathcal{A}_W with

respect to the composition of morphisms in \mathcal{A}_W .

Proof: Since $f_\omega(\omega') = \omega\omega'$, it follows that $f_{\omega_1} f_{\omega_2} = f_{\omega_1 \omega_2}$.

3.4 From Prop. 3.3 it follows that for any object A of \mathcal{A}_W , the set

$H_{M_W}(A)$ enjoys a structure of a transition system with input W by

combining $H_{M_W}(A)$ with $W \xrightarrow{R} \text{End}_{\mathcal{A}_W}(M_W)$.

Formally, we define a functor $\text{Mor} : \mathcal{A}_W \rightarrow \mathcal{A}_W$, where for

any object A of \mathcal{A}_W we define $\text{Mor}(A)$ by:

$$S(\text{Mor}(A)) = H_{M_W}(A),$$

$$f_s \cdot \omega = f_s f_\omega = f_{s \cdot \omega} \text{ for any } M_W \xrightarrow{f_s} A \text{ and } \omega \in W.$$

For any morphism $A \xrightarrow{g} B$ we define $\text{Mor}(A) \xrightarrow{\text{Mor}(g)} \text{Mor}(B)$

by $\text{Mor}(g) = H_{M_W}(g)$.

An immediate verification shows that $\text{Mor}(A)$ is an object of

\mathcal{A}_W , and that $H_{M_W}(g)$ determines in fact a morphism of \mathcal{A}_W . Furthermore,

it follows directly from the fact that H_{M_W} is a functor that $\text{Mor} : \mathcal{A}_W \rightarrow \mathcal{A}_W$

is also a functor. Likewise, the transformation $\rho: S \rightarrow H_{M_W}$

determines directly a transformation $\rho_{A_W}: I \rightarrow \text{Mor}$ from the identity

functor of A_W to Mor , and we have:

3.4.1 THEOREM: The transformation $\rho_{A_W}: I \rightarrow \text{Mor}$ is a natural

equivalence of functors.

3.4.2 Intuitively speaking, the functor Mor constructs the "internal structure" of any object A of A_W from a part of the category A_W which lies around M_W and between M_W and A . Hence it is intuitively clear, that if M_W can be recognized in A_W (up to an isomorphism) by means of some categorical predicate, then the "internal structure" of any object can be reconstructed "categorically," and therefore any property of the transition systems with input W can be determined "categorically" as well.

3.5 LEMMA: If W is a unit-commutative monoid (i.e., if $uv = 1_W$ in W then $vu = 1_W$) or a finite monoid then every epic morphism $A \xrightarrow{e} M_W$ of A_W , where A is monogenic, is an isomorphism.

Proof: If W is a finite monoid then the cardinality of the set of states of any monogenic transition system with input W cannot exceed the cardinality of W . Hence $e : S(A) \rightarrow W$ must be bijective.

If W is a unit-commutative monoid and A is generated by $\{s_0\}$, then $\{e(s_0)\}$ must generate M_W , that is, $e(s_0)v = 1_W$ for some $v \in W$, and therefore $ve(s_0) = 1_W$.

Assume that $e(s_0 \cdot \omega_1) = e(s_0 \cdot \omega_2)$ for some $\omega_1, \omega_2 \in W$, then we have $\omega_1 = ve(s_0)\omega_1 = ve(s_0 \cdot \omega_1) = ve(s_0 \cdot \omega_2) = ve(s_0)\omega_2 = \omega_2$, and therefore $s \cdot \omega_1 = s \cdot \omega_2$, which shows that e is also injective.

3.5.1 COROLLARY: If W is a unit-commutative monoid or a finite monoid, then for any automorphism F of A_W , $F(M_W)$ is isomorphic to M_W .

Proof: From lemma 3.5 it follows that an object M of

\mathcal{A}_W is isomorphic to M_W iff

(i) M is monogenic, and

(ii) for any monogenic object A of \mathcal{A}_W there exists an epic morphism $M \xrightarrow{e} A$ of \mathcal{A}_W .

Since these properties of morphisms and objects of \mathcal{A}_W are preserved under the automorphisms of \mathcal{A}_W , the corollary follows.

3.5.2 Note that the class of unit-commutative monoids is broad enough to cover all the classes of monoids which are employed in automata theory. For example, the left-cancellative and the right-cancellative monoids are all unit-commutative. Hence the free monoids and the groups are unit-commutative. Note also that the cartesian products of unit-commutative monoids are unit-commutative, and therefore we can apply our results to multi-input transition systems as well.

4. CATEGORICAL PREDICATES AND TRANSPARENT CATEGORIES

4.1 A subcategory \mathcal{D} of \mathcal{C} is said to be very full iff for any morphism h of \mathcal{D} and for any morphisms f and g of \mathcal{C} such that $fg = h$, $fh = g$ or $hf = g$ holds in \mathcal{C} it follows that f and g belong to \mathcal{D} .

A functor $T : \mathcal{D} \rightarrow \mathcal{C}$ is said to be a very full embedding iff T is an embedding and the image of T is a very full subcategory of \mathcal{C} .

Let \mathcal{D} be a category and D a class of morphisms of \mathcal{D} , we denote by $(\mathcal{D}, D, \mathcal{C})$ the class of all the images of the morphisms in D under any very full embedding $T : \mathcal{D} \rightarrow \mathcal{C}$. That is, $f \in (\mathcal{D}, D, \mathcal{C})$ iff there are a morphism $d \in D$ and a very full embedding $T : \mathcal{D} \rightarrow \mathcal{C}$ such that $f = T(d)$.

A class K of morphisms of \mathcal{C} is said to be categorical (in \mathcal{C}) iff there is a category \mathcal{D} such that $K = (\mathcal{D}, D, \mathcal{C})$ for some class D of morphisms of \mathcal{D} .

4.2 PROPOSITION: A class K is categorical in \mathcal{C} iff it is closed under all the automorphisms of \mathcal{C} .

Proof: Since for any very full embedding functor $T : \mathcal{D} \rightarrow \mathcal{C}$ and for any automorphism F of \mathcal{C} , $F \circ T : \mathcal{D} \rightarrow \mathcal{C}$ is also a very full embedding, it follows that every categorical class in \mathcal{C} is closed under all the automorphisms of \mathcal{C} .

On the other hand, let K be a class of morphisms of \mathcal{C} which is closed under all automorphisms of \mathcal{C} . Denote by $\mathcal{D}(K)$ the minimal very full subcategory of \mathcal{C} which includes K , then $K = (\mathcal{D}(K), K, \mathcal{C})$.

In order to see this, let $T : \mathcal{D}(K) \rightarrow \mathcal{C}$ be any very full embedding and define $F_T : \mathcal{C} \rightarrow \mathcal{C}$ by

$$F_T(f) = \begin{cases} T(f) & \text{if } f \in \mathcal{D}(K), \\ f & \text{otherwise.} \end{cases}$$

Since $\mathcal{D}(K)$ is a very full subcategory of \mathcal{C} , F_T maps $\mathcal{D}(K)$ into itself, and because T is a very full embedding, F_T maps $\mathcal{D}(K)$ onto itself in an injective manner. Furthermore, F_T must be a functor and it has an inverse, hence it is an automorphism of \mathcal{C} .

Now, since K is closed under automorphisms it follows that $(\mathcal{D}(K), K, \mathcal{C}) \subseteq K$, and since clearly $K \subseteq (\mathcal{D}(K), K, \mathcal{C})$ we have the

the desired equality.

4.2.1 COROLLARY: A class K is categorical in \mathcal{C} iff $K = (\mathcal{D}, \mathcal{D}, \mathcal{C})$

for some very full subcategory \mathcal{D} of \mathcal{C} .

4.2.2 COROLLARY: Let \mathcal{D} be any category and \mathcal{D} a class of some morphisms of \mathcal{D} , then the class of all values of the morphisms in \mathcal{D} under all embedding functors $\mathcal{D} \rightarrow \mathcal{C}$ is categorical in \mathcal{C} .

4.2.3 COROLLARY: The class of all values of the morphisms in \mathcal{D} under all functors $\mathcal{D} \rightarrow \mathcal{C}$ is categorical in \mathcal{C} .

4.2.4 Note that we cannot dispense with the requirement of employing very full embeddings in the definition of the categorical classes in any arbitrary category. For example in the category \mathcal{N} of natural numbers where the morphisms represent the natural partial order of natural numbers, every set of morphisms is categorical.

However, the categorical classes achieved by means of 4.2.2 or 4.2.3 are always infinite or empty.

4.3.1 A class of morphisms of \mathcal{C} is said to be natural iff it is closed under all those automorphisms of \mathcal{C} which are naturally equivalent to $I_{\mathcal{C}}$, the identity functor of \mathcal{C} .

Obviously, by Prop. 4.2 we have that every categorical class is natural. Note that a class of identity morphisms of \mathcal{C} which is closed under the isomorphisms within \mathcal{C} is always natural.

4.3.2 A category \mathcal{C} is said to be transparent if all the natural classes in \mathcal{C} are categorical.

4.4 Obviously, if all the automorphisms of \mathcal{C} are naturally equivalent (i.e., to $I_{\mathcal{C}}$) then \mathcal{C} is transparent.

Let us call a category \mathcal{C} autotrivial iff all the automorphisms of \mathcal{C} are naturally equivalent.

It is not known whether all transparent categories are autotrivial. All "natural" categories of mathematical systems that are known to be transparent are in fact autotrivial as well.

The equivalence between the notion of transparent categories and that of autotrivial categories, in a special case, takes the form of the following problem in group theory:

Do all groups whose automorphisms are all (conjugate) class preserving have only inner automorphisms?

Any example of a group all of whose automorphisms are class preserving and which has an outer automorphism, yields a transparent category (with a single object and all its morphisms are invertible and in one-one correspondence with the elements of the group) which is not autotrivial.

5. THE DISTINGUISHABILITY OF M_W AND THE TRANSPARENCY OF A_W .

5.1 The features exhibited by M_W in A_W are quite common in "natural" categories. As we shall see presently, they provide a reduction of the autotriviality of categories to the categoricity of certain classes of identity morphisms.

An object M of a category \mathcal{C} is said to be a generator of \mathcal{C} iff $H_M : \mathcal{C} \rightarrow \mathcal{S}$ (where $H_M(A) = \text{Hom}_{\mathcal{C}}(M, A)$) is an embedding. In this case, the values of H_M form a subcategory $H_M(\mathcal{C})$ of \mathcal{S} . In particular, M is said to be a faithful generator of \mathcal{C} iff there exists a functor $R_M : H_M(\mathcal{C}) \rightarrow \mathcal{C}$ such that $R_M \circ H_M : \mathcal{C} \rightarrow \mathcal{C}$ is naturally equivalent to $I_{\mathcal{C}}$.

5.1.1 EXAMPLES: The additive group Z of integers is a faithful generator of both the category of abelian groups and the category of all groups (Freyd 1964). The single-element set U is a faithful generator of \mathcal{S} the category of sets; in fact H_U is already naturally equivalent to the

identity functor of \mathcal{S} .

From our results in Chapter 3 we know that M_W is a faithful generator of \mathcal{A}_W .

5.2 LEMMA: For any object A of \mathcal{C} and any automorphism F of \mathcal{C} with inverse G , $H_A \circ F$ is naturally equivalent to $H_{G(A)}$

Proof: We shall prove a stronger result; namely, for any two objects A and B of \mathcal{C} and any automorphism F of \mathcal{C} with inverse G , there exists a bijection

$$\Phi(A, B) : \text{Hom}_{\mathcal{C}}(G(A), B) \longrightarrow \text{Hom}_{\mathcal{C}}(A, F(B))$$

which is natural in both A and B .

Put differently, F and its inverse are adjoint. By Kan's characterization of adjoint functors (Kan 1958, MacLane 1965) the following is a proof that F and G are adjoint (to each other!):

Let us denote by e_C the identity morphism of an arbitrary object C of \mathcal{C} .

(i) Every morphism $A \xrightarrow{f} F(B)$ can be factored as $f = F(h) \cdot e_A$

for some $G(A) \xrightarrow{h} B$ (i.e., $h = G(f)$).

(ii) If $F(h_1) \cdot e_A = F(h_2) \cdot e_A$ then obviously $h_1 = h_2$.

5.3 Let M be a faithful generator of \mathcal{C} . From Lemma 5.2 we know

that $H_M \circ F$ is naturally equivalent to $H_{G(M)}$ for any automorphism F of \mathcal{C} with inverse G . Hence F , which is naturally equivalent to $R_M \circ H_M \circ F$,

is naturally equivalent to $R_M \circ H_{G(M)}$. If $G(M)$ is isomorphic to M ,

then H_M is naturally equivalent to $H_{G(M)}$, and therefore F , which is

naturally equivalent to $R_M \circ H_{G(M)}$, is naturally equivalent to $I_{\mathcal{C}}$.

5.3.1 An object A of \mathcal{C} is said to be distinguishable (in \mathcal{C}) iff for

any automorphism F of \mathcal{C} , $F(A)$ is isomorphic to A . Put differently,

A is distinguishable in \mathcal{C} iff the natural class of all identity

morphisms of the objects of \mathcal{C} , which are isomorphic to A in \mathcal{C} , is

categorical.

Thus we have proved:

5.3.2 THEOREM: A category \mathcal{C} with a faithful generator M is

autotrivial iff M is distinguishable in \mathcal{C} .

5.3.3 COROLLARY: A category \mathcal{C} with a faithful generator is autotrivial

iff it is transparent.

5.4 Since M_W is a faithful generator of \mathcal{A}_W , \mathcal{A}_W is transparent iff

it is autotrivial. Furthermore \mathcal{A}_W is autotrivial iff M_W is dis-

tinguishable in \mathcal{A}_W . By 3.5.1 we know that if W is finite or unit-

commutative then M_W is distinguishable in \mathcal{A}_W . Thus we have:

5.4.1 THEOREM: If W is a finite monoid or a unit commutative

monoid then \mathcal{A}_W is autotrivial (and therefore transparent).

6. DISCUSSION AND OPEN PROBLEMS

6.1 Our result as expressed by Theorem 5.4.1 implies that, for a very broad class of monoids, the categorical study of a domain of all transition systems with input monoid of this class, is equivalent in principle to the "complete" study (or the "inside" study) of these systems. However, only experience may show us that in fact there is a psychological advantage to the categorical approach in the study of these systems

6.2 If G is a group then A_G is the category of all representations of G as operating on sets. Since every group is in particular a unit-commutative monoid, we have that the categorical study of the representations of a fixed arbitrary group G is sufficient in principle for producing all the algebraic properties of the representations of G .

6.3 Our results so far, give rise to some general problems that deserve attention. For example, is it true that for any monoid W , M_W is distinguishable in A_W ?

More generally, what additional properties on faithful generators of categories, if any at all, are necessary in order to insure that they are distinguishable? In particular, is it true that every projective faithful generator is distinguishable?

6.4 Important and much more interesting categories of transition systems are those of finite-state transition systems (i.e., transition systems whose sets of states are finite). If the input monoid W is also finite then our results remain valid since M_W is also finite. If however W is infinite, as it is the case in the ordinary theory of finite automata (where the input monoids are finitely generated free monoids) then M_W is no longer applicable.

6.5 Another interesting restriction of A_W is to abelian transition

systems. A transition system A is said to be abelian iff

$s \cdot \omega_1 \omega_2 = s \cdot \omega_2 \omega_1$ holds for all $s \in S(A)$ and $\omega_1, \omega_2 \in W$.

For any arbitrary monoid W there exists a homomorphism of monoids $W \xrightarrow{ab} W^{ab}$ where W^{ab} is an abelian monoid with the following universality property : any homomorphism of W into an abelian monoid factors uniquely through ab . The direct construction of W^{ab} must be evident.

Denote by M_W^{ab} the following object of A_W :

$$S(M_W^{ab}) = W^{ab},$$

$$ab(\omega_1) \cdot \omega_2 = ab(\omega_1 \omega_2) .$$

Obviously, M_W^{ab} is abelian. If we denote by A_W^{ab} the full subcategory of A_W of abelian objects, one can easily follow the example of M_W in A_W and show that M_W^{ab} is a fait.fu' generator of A_W^{ab} . Furthermore, for any arbitrary monoid W , M_W^{ab} is distinguishable by the same properties which distinguish M_W in A_W in the case of unit-commutative input monoid W (cf. 3.5.1), hence for any arbitrary monoid W , A_W^{ab} is transparent and autotrivial.

An equivalent proof of these properties of A_W^{ab} follows directly from the fact that A_W^{ab} is a category isomorphic to $A_{W^{ab}}$.

BIBLIOGRAPHY

Freyd, P., "Abelian Categories: An Introduction to the Theory of Functors," Harper's Series in Modern Mathematics, New York (1964).

Give'on, Y., "Toward a Homological Algebra of Automata I: The Representation and Completeness Theorem for Categories of Abstract Automata," Technical Report, Department of Communication Sciences, ORA, The University of Michigan (1964).

Give'on, Y., "Toward a Homological Algebra of Automata," Technical Report, Department of Communication Sciences, ORA, The University of Michigan (1965).

Hartmanis, J. and Stearns, R. E., "Pair Algebra and Automata Theory," Inform. Control, 7, 485 - 507 (1964).

Kan, D. M., "Adjoint Functors," Trans. Amer. Math. Soc., 87, 295 - 329 (1958).

MacLane, S., "Categorical Algebra," Bull. Amer. Math. Soc., 71, 40 - 106 (1965).

Zeiger, H. P., "Loop-Free Synthesis of Finite State Machines," thesis, MIT (1964).

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) Logic of Computers Group The University of Michigan Ann Arbor, Michigan 48104		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP
3. REPORT TITLE TRANSPARENT CATEGORIES AND CATEGORIES OF TRANSITION SYSTEMS		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (Last name, first name, initial) Give'on, Yehoshafat		
6. REPORT DATE May, 1965	7a. TOTAL NO. OF PAGES 35	7b. NO. OF REFS 7
8a. CONTRACT OR GRANT NO. Nonr 1224 (21)	9a. ORIGINATOR'S REPORT NUMBER(S) 03105-38-T	
a. PROJECT NO.		
c.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		
10. AVAILABILITY/LIMITATION NOTICES Qualified requesters may obtain copies of this report from DDC.		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY Office of Naval Research Department of the Navy Washington 25, D.C.	
13. ABSTRACT / This paper presents a study of the homomorphisms of transition systems, which are the main tool employed in recent studies in algebraic automata theory. The problem of determining the extent of the information that can be retrieved from the algebra of these homomorphisms is defined and discussed. Naturally, this study is allied with categorical algebra. A general result which is applicable to many "natural" categories of mathematical systems is derived. This result implies the completeness of the categorical algebra study of transition systems with a fixed input monoid, provided that the input monoid belongs to a very broad class of monoids which includes all the types of monoids which are employed in automata theory.		

DD FORM 1473
1 JAN 64

Unclassified

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
algebraic automata theory categorical algebra						

INSTRUCTIONS

1. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. REPORT DATE: Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. NUMBER OF REFERENCES: Enter the total number of references cited in the report.

8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.

12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.